

The Physiological Effects of Hyperthermia¹

Haim I. Bicher, M.D., Ph.D.

In the application of hyperthermia to control cancer, physiological factors such as pH, pO₂, and blood flow must be documented *in vivo* at normal and elevated temperatures for both the tumor and its normal host tissue. The present symposium was arranged to gather recent research done in this area and to expose the problems inherent in these efforts.

INDEX TERMS: Blood, flow dynamics • Hyperthermia • Oxygen

Radiology 137:511-513, November 1980

HYPERTHERMIA is currently being employed in clinical trials to control cancer. The protocols in these trials employ hyperthermia either as the only treatment modality or in combination with other accepted modalities such as radiation therapy and chemotherapy.

The current interest in hyperthermia is prompted mainly by some very interesting radiobiological observations made at the cellular level. *In vitro* studies (1-4) demonstrate that heat-induced lethal damage to cells in culture follows a quasithreshold exponential response similar in shape to cell survival curves obtained for ionizing radiation exposure or treatment with some cytotoxic agents. Different cell lines exhibit widely different thermal sensitivities (2, 3). Heat survival curves at 45.5°C for aerobic and hypoxic cells using CHO cells show that hypoxic cells are more heat sensitive than well oxygenated cells (5). Similar observations were made by Schulman and Hall (6) for V79 Chinese hamster cells and by Kim *et al.* (7) for HeLa cells. Assuming that the W138VA 13/2 RA cells model the malignant cells of a human tumor and W138 cells model the associated normal tissue, Kase and Hahn (8) observed that the transformed cells are more readily destroyed by hyperthermia. Others (9-13) have also reported that tumor cells are more thermosensitive than normal cells. Just as in the case of ionizing radiation, the response to hyperthermia is also dependent on the phase of the cell cycle during exposure. However, mid-to-late S-phase cells, which are the most resistant to ionizing radiation, are the most sensitive to hyperthermia (4, 12, 14). Gerweck (15, 16) has clearly demonstrated that the lethal response of CHO cells to hyperthermia (41-44°C) increases as pH is reduced from 7.6 to 6.7. Hahn (17) has shown that when cells are heated in a salt solution lacking nutrients instead of in a complete medium, thermal sensitivity increases greatly. Another aspect of hyperthermia which may have a great impact on its success is the resistance of cells to subsequent heat treatments. The phenomenon is termed thermotolerance. From studies with mammalian cells in

culture (18, 19) it was observed that there is an increase in D₀ of a subsequent hyperthermia survival curve. Thermotolerance is induced during a 3-5 hour incubation period at near physiological temperatures after heating for a short time at temperatures greater than 43°C (20, 21).

In vitro studies (7, 10, 22, 23) also have demonstrated that hyperthermia combined with ionizing radiation has a synergistic cell killing effect. The importance of the sequence in which heat and radiation are applied has been studied in detail by Sapareto *et al.* (22). The synergistic effect is most pronounced for radioresistant S-phase cells (22, 24). There are no *in vitro* data indicating differential sensitization of cancer cells as opposed to normal tissue cells.

In the recent literature (26-38), several investigators have achieved local tumor control using hyperthermia either alone or in combination with irradiation. Crile (26) studied the effectiveness of hyperthermia and the dependence of cure on the temperature achieved and the length of the treatment period on mouse sarcoma. He also demonstrated the effectiveness of the combination of heat and radiation in the treatment of cancer of the rectum (27). Hartman and Crile (28) applied hyperthermia in the treatment of human osteogenic sarcoma with good results. Cavaliere *et al.* (29) and Morica *et al.* (30) documented the effect of heat on tumor cells in rats and the sparing of normal tissues. Other investigators (31, 32) had similar data showing normal tissue sparing with the combined modalities. Recent clinical reports (33-38) clearly demonstrate that hyperthermia, particularly in combination with irradiation, can prove to be a very useful modality in the treatment of cancer.

In vivo results do not always agree with the *in vitro* observations made under controlled conditions. This is probably due to the interaction of various physiological factors not present in the *in vitro* systems. An interesting example is the conflict between the *in vitro* results of Sapareto *et al.* (23) and the *in vivo* results of Law *et al.* (25)

¹ From The Department of Therapeutic Radiology, Division of Radiation Biology and Physics (H.I.B., Division Head), Henry Ford Hospital, Detroit, MI 48202. Submitted for publication 5 April 1980; accepted 1 July 1980.

Presented at the Hyperthermia Microcirculation Symposium conducted by the Radiation Therapy Oncology Group under the sponsorship of the National Institutes of Health, Albuquerque, NM, Jan 1980. See also the following articles by Eddy (p. 515) and Bicher (p. 523) in this issue. The remainder of the Symposium will be published in the December 1980 issue of RADILOGY. shan

when combining hyperthermia and irradiation. The *in vitro* results indicate that, with heating time adjusted to achieve the same killing from heat alone at various temperatures, a decrease in temperature increases the relative killing from the interaction of radiation and heat. This is in contradiction to the *in vivo* results (25) which indicate that the opposite should occur.

In order to fully make use of *in vitro* observations made under controlled conditions in the application of hyperthermia to cancer control, the physiological factors such as pH, oxygen consumption, nutrients, and blood flow of both tumor and normal host tissue should be measured *in vivo*. Furthermore, these factors should be documented not only at normal temperatures but also under hyperthermic conditions. Under certain conditions a selective destruction of tumor tissue might be possible with normal tissue sparing. For example, several studies (39–42) indicate that pH of fluid in human and rodent solid tumors is lower than the normal tissue pH of 7.4. Paramount among the other factors which may change and subsequently influence the response of cells or tissues to supranormal temperatures are the vascular changes, blood flow response and the net result of this on tissue oxygenation. The last factor may change the effect of both hyperthermia and radiation therapy when used in combination. Insufficient information is available on the temperature modifications of tumor blood supply, tumor oxygen tension and consumption and respiratory gas exchange by malignant cells *in vivo*.

This symposium on the physiological effects of hyperthermia was arranged in order to assemble recent research on the physiological response of normal and tumor tissue to hyperthermia.

Haim I. Bicher, M.D., Ph.D.
Staff Radiotherapist
Head, Division of Radiobiology and Physics
Henry Ford Hospital
Detroit, MI 48202

REFERENCES

- Dewey WC, Hopwood LE, Sapareto SA, et al: Cellular responses to combinations of hyperthermia and radiation. *Radiology* **123**: 463–474, May 1977
- Harris M: Temperature-resistant variants in clonal populations of pig kidney cells. *Exp Cell Res* **46**:301–314, May 1967
- Westra A, Dewey WC: Variation in sensitivity to heat shock during the cell-cycle of Chinese hamster cells *in vitro*. *Int J Radiat Biol* **19**:467–477, May 1971
- Palzer RJ, Heidelberger C: Studies on quantitative biology of hyperthermic killing of HeLa cells. *Cancer Res* **33**:415–421, Feb 1973
- Gerweck LE, Gillette EL, Dewey WC: Killing of Chinese hamster cells *in vitro* by heating under hypoxic or aerobic conditions (letter). *Eur J Cancer* **10**:691–693, Oct 1974
- Schulman N, Hall EJ: Hyperthermia: its effect on proliferative and plateau phase cell cultures. *Radiology* **113**:209–211, Oct 1974
- Kim SH, Kim JH, Hahn EW: Enhanced killing of hypoxic tumour cells by hyperthermia (letter). *Br J Radiol* **48**:872–874, Oct 1975
- Kase K, Hahn GM: Differential heat response of normal and transformed human cells in tissue culture. *Nature* **255**:228–230, 15 May 1975
- Bronk VV: Thermal potentiation of mammalian cell killing. Clues for understanding and potential for tumor therapy. (In) Lett JT, Adler H, Felle M, eds: *Advances in Radiation Biology*. New York, Academic Press, Vol VI, 1976, pp 267–323
- Thrall DE, Gerweck LE, Gillette EL, et al: Responses of cells *in vitro* and tissues *in vivo* to hyperthermia and x-irradiation. (In) Lett JT, Adler H, Felle M, eds: *Advances in Radiation Biology*. New York, Academic Press, Vol. VI, 1976, pp 211–227
- Har-Kedar I, Bleehen NM: Experimental and clinical aspects of hyperthermia applied to the treatment of cancer with special reference to the role of ultrasonic and microwave heating. (In) Lett JT, Adler H, Felle M, eds: *Advances in Radiation Biology*. New York, Academic Press, Vol VI, 1976, pp 228–266
- Palzer RJ, Heidelberger C: Influence of drugs and synchrony on the hyperthermic killing of HeLa cells. *Cancer Res* **33**:422–427, Feb 1973
- Chen TT, Heidelberger C: Quantitative studies on the malignant transformation of mouse prostate cells by carcinogenic hydrocarbons *in vitro*. *Int J Cancer* **4**:166–178, 15 Mar 1969
- Kim SH, Kim JH, Hahn EW: The enhanced killing of irradiated HeLa cells in synchronous culture by hyperthermia. *Radiat Res* **66**:337–345, May 1976
- Gerweck LE, Rottinger E: Enhancement of mammalian cell sensitivity to hyperthermia by pH alteration. *Radiat Res* **67**: 508–511, Sep 1976
- Gerweck LE: Modification of cell lethality at elevated temperatures. The pH effect. *Radiat Res* **70**:224–235, Apr 1977
- Hahn GM: Metabolic aspects of the role of hyperthermia in mammalian cell inactivation and their possible relevance to cancer treatment. *Cancer Res* **34**: 3117–3123, Nov 1974
- Gerner EW, Schneider MJ: Induced thermal resistance in HeLa cells. *Nature* **256**:500–502, 7 Aug 1975
- Henle KJ, Leeper DB: Interaction of hyperthermia and radiation in CHO cells: recovery kinetics. *Radiat Res* **66**:505–518, Jun 1976
- Harisiadis L, Sung DI, Hall EJ: Thermal tolerance and repair of thermal damage by cultured cells. *Radiology* **123**:505–509, May 1977
- Henle KJ, Karamuz JE, Leeper DB: Induction of thermotolerance in Chinese hamster ovary cells by high (45°C) or low (40°C) hyperthermia. *Cancer Res* **38**:570–574, Mar 1978
- Sapareto SA, Hopwood LE, Dewey WC: Combined effects of x-irradiation and hyperthermia on CHO cells for various temperatures and orders of application. *Radiat Res* **73**:221–233, Feb 1978
- Sapareto SA, Raaphorst PG, Dewey WC: Cell killing and the sequencing of hyperthermia and radiation. *Int J Radiat Oncol Biol Phys* **5**:343–347, Mar 1979
- Gerweck LE, Gillette EL, Dewey WC: Effect of heat and radiation on synchronous Chinese hamster cells: killing and repair. *Radiat Res* **64**:611–623, Dec 1975
- Law MP, Ahier RG, Field SB: The response of the mouse ear to heat applied alone or combined with x-rays. *Br J Radiol* **51**: 132–138, Feb 1978
- Crile G Jr: The effects of heat and radiation on cancer implanted on the feet of mice. *Cancer Res* **23**:372–380, Mar 1963
- Crile G Jr: Heat as an adjunct to the treatment of cancer. *Cleveland Clin Q* **28**:75–89, Apr 1961
- Hartman JT, Crile G Jr: Heat treatment of osteogenic sarcoma. Report of five cases. *Clin Orthop* **61**:269–276, Nov–Dec 1968
- Cavaliere R, Ciocatto EC, Giovanella BC, et al: Selective heat sensitivity of cancer cells. *Cancer* **20**:1351–1381, Sep 1967
- Moricca G, Cavaliere R, Caputo A, et al: Hyperthermic treatment of tumours. Experimental and clinical applications. *Recent Results Cancer Res* **59**:112–152, 1977
- Robinson JE, Wizenberg MJ, McCready WA: Radiation and hyperthermal response of normal tissue *in situ*. *Radiology* **113**: 195–198, Oct 1974
- Overgaard K, Overgaard J: Investigations on the possibility of

- a thermic tumour therapy. I. Short-wave treatment of a transplanted isologous mouse mammary carcinoma. *Eur J Cancer* **8**:65-78, Feb 1972
33. LeVeen HH, Wapnick S, Piccone V, et al: Tumor eradication by radiofrequency therapy. *JAMA* **235**:2198-2200, 17 May 1976
 34. Hornback NB, Shupe RE, Shidnia H, et al: Preliminary clinical results of combined 433 megaHertz microwave therapy and radiation therapy on patients with advanced cancer. *Cancer* **40**: 2854-2863, Dec 1977
 35. Johnson RJ, Sandhu TS, Hetzel FW, et al: A pilot study to investigate the therapeutic ratio of hyperthermia (41.5-42.0°C) and radiation. *Int J Radiat Oncol Biol Phys* **5**:549-555, Jun 1979
 36. Kim JH, Hahn EW, Benjamin FJ: Treatment of superficial cancers by combination hyperthermia and radiation therapy. *Clin Bull* **9**:13-16, 1979
 37. Storm FK, Harrison WH, Elliott RS, et al: Normal tissue and solid tumor effects of hyperthermia in animal models and clinical trials. *Cancer Res* **39**:2245-2251, Jun 1979
 38. Bicher HI, Sandhu TS, Hetzel FW: Hyperthermia as an adjuvant to radiation: Proposal for an effective fractionation regime. *Int J Radiat Oncol Biol Phys* **6**:867-870, Jul 1980
 39. Meyer KA, Kammerling EM, Antman L, et al: pH studies of malignant tissues in human beings. *Cancer Res* **8**:513-518, 1948
 40. Naeslund J, Swenson KE: Investigations on the pH of malignant tumours in mice and humans after administration of glucose. *Acta Obstet Gynecol Scand* **32**:359-367, 1953
 41. Eden M, Haines B, Kahler H: The pH of rat tumors measured in vivo. *J Natl Cancer Inst* **16**:541-556, Oct 1955
 42. Gullino PM, Grantham FH, Smith SH, et al: Modifications of the acid-base status of the internal milieu of tumors. *J Natl Cancer Inst* **34**:857-869, Jun 1965